arXiv:1302.5451v1 [math.GR] 21 Feb 2013

Strong accessibility for finitely presented groups

Larsen Louder & Nicholas Touikan

February 25, 2013

Abstract

A hierarchy of a group is a tree of groups obtained by iteratively passing to vertex groups of graphs of groups decompositions. We show that given a slender hyperbolic hierarchy $\mathcal{H}$ of a finitely presented group H there is some level N and some constant C such that each group in $\mathcal{H}$ at level $n \geq N$ has a finite slender hierarchy of height at most C such that terminal leaves are either slender or elliptic in the original hierarchy $\mathcal{H}$.

As a corollary we deduce that a finitely presented group which contains no slender groups which act dihedrally on the line has a hierarchy whose terminal groups are either slender or don't split over a slender subgroup.

1 Introduction

A group G is said to be *accessible* over a family of subgroups $\mathcal{C}$ if there is an upper bound to the size of reduced graphs of groups decompositions of G with edge groups in $\mathcal{C}$. The classic theorem is due to Grushko: If $G = A * B$ is a nontrivial decomposition of G as a free product then $\mathrm{rk}(A) + \mathrm{rk}(B) = \mathrm{rk}(G)$, which implies that there is an upper bound to the size of reduced graphs of groups decompositions of a given finitely generated group G over trivial edge groups.

Finitely generated groups are not accessible over the class of small subgroups, as Dunwoody and Bestvina/Feighn have produced counterexamples (with finite and small edge groups, respectively) [Dun93, BF91b]. Finitely presented groups, on the other hand, are accessible over the class of small subgroups [Dun85, BF91a].

It is natural to try to repeat this, and find conditions which guarantee that the process of iteratively passing to vertex groups of graphs of groups decompositions terminates in finitely many steps ("strong accessibility"). For finitely generated groups this fails for both silly (there is an HNN extension $\mathbb{F}_2 \cong \mathbb{F}_2 *_{\mathbb{Z}}$) and mysterious reasons: there is a finitely generated one-ended group G such that $G = G *_C G$, with C infinite cyclic [DJ98]. Delzant and Potyagailo claim in [DP01] that finitely presented groups admit finite hierarchies over "elementary families", but unfortunately the proof of [DP01, Lemma 4.10] is not correct. See Example 1.

We know of two non-artificial classes of groups which have hierarchies that must terminate. A Haken hierarchy of a three-manifold gives a finite hierarchy in this sense: an incompressible two-sided surface in a three-manifold corresponds to a splitting of its fundamental group over the fundamental group of the surface. Sela and, independently, Kharlampovich, Myasnikov and Remeslennikov have shown that the hierarchy of a limit group obtained by alternatingly passing to vertex groups of the Grushko and abelian JSJ decomposition is finite.

A group is *small* if it doesn't contain a non-abelian free subgroup. An action of a small group on a tree is either *elliptic* (fixes a point in the tree) *hyperbolic* (has an axis and acts by translations) *dihedral* (has an axis and acts dihedrally) or *parabolic* (fixes a point in the boundary but has no axis) [BF91a, p. 453]. A group is *slender* if all its subgroups are finitely generated. An action of a slender group on a tree cannot be parabolic.

Definition 1.1 (Hierarchy)**.** A *hierarchy* for a group H is a rooted tree of groups $\mathcal{H}$, with H at the root, such that the descendants of a group $L \in \mathcal{H}$ are the vertex groups of a nontrivial graph of groups decomposition Δ_L of L.

A hierarchy is *slender* if all graphs of groups decompositions are over slender edge groups. A slender hierarchy is *hyperbolic* if for all $L \in \mathcal{H}$, if $E < L$ is slender, then the action of E on T_{Δ_L}, the tree associated to Δ_L, is either elliptic or hyperbolic.

Denote the groups at level n of the hierarchy by $\mathcal{H}^n$. A subgroup $V < L \in \mathcal{H}^n$ is *$\mathcal{H}$–elliptic* if there is a chain $L = L_n > L_{n+1} > L_{n+1} > \dots$ such that $L_i \in \mathcal{H}^i$ is a vertex group of $\Delta_{L_{i-1}}$ and elements $h_i \in L_i$ such that

$$V < L_n \cap L_{n+1}^{h_n} \cap L_{n+2}^{h_{n+1}h_n} \cap L_{n+3}^{h_{n+2}h_{n+1}h_n} \cap \cdots$$

Recall that a group is almost finitely presented if it acts freely and cocompactly on a connected simplicial complex X with $H^1(X, \mathbb{Z}_2) = 0$ [Dun85]. We use a slightly less restrictive notion of almost finitely presented.

Definition 1.2 ($\mathcal{H}$–almost finitely presented)**.** Let H be a finitely generated group and $\mathcal{E}$ a family of subgroups of H. We say that H is *almost finitely presented relative to $\mathcal{E}$* if H acts cocompactly on a connected triangular complex X such that $H^1(X, \mathbb{Z}_2) = 0$ and cell stabilizers in X are either slender or conjugate into $\mathcal{E}$.

Let H be finitely generated and let $\mathcal{H}$ be a hierarchy of H. Call H *$\mathcal{H}$–almost finitely presented* H is almost finitely presented relative to the family of $\mathcal{H}$–elliptic subgroups.

Theorem 1.3 (Main theorem)**.** *Let H be a finitely generated group and let $\mathcal{H}$ be a slender hyperbolic hierarchy of H. If H is $\mathcal{H}$–almost finitely presented then for some N, there is a constant C such that each $L \in \mathcal{H}^m$, $m \geq N$, has a finite hierarchy $\mathcal{X}_L$ of height at most C whose terminal groups are either $\mathcal{H}$–elliptic or slender.*

Corollary 1.4. *Let H be finitely presented and suppose H contains no slender groups with an infinite dihedral ($\mathbb{Z}_2 * \mathbb{Z}_2$) quotient. Let $\mathcal{H}$ be the hierarchy such that Δ_L is a slender* JSJ *decomposition (See [FP06].) of L, and such that if $L \in \mathcal{H}$ is slender then we assume L is terminal. Then $\mathcal{H}$ is finite.*

Let H be torsion-free relatively hyperbolic. The hierarchy such that Δ_L is either the Grushko (if L is freely decomposable) or JSJ *decomposition (if L is freely indecomposable) relative to the collection of peripheral subgroups, is finite. If H is toral relatively hyperbolic the same holds but for the full abelian* JSJ *decomposition.*

Disallowing slender groups with dihedral quotients is undesirable.

Question 1. *Does Corollary 1.4 hold for the hierarchy obtained by passing to the* JSJ *decomposition relative to slender subgroups acting dihedrally in the tree associated to the slender* JSJ *decomposition?*

Acknowledgments

We would like to thank Thomas Delzant and Leonid Potyagailo for some interesting conversations. Thanks to Daniel Groves for help with an earlier version of the paper, as well as to Peter Haïssinsky and Matias Carrasco Piaggio for encouragement, and to Thierry Coulbois for bringing our attention to [Gui05].

The first author was supported in part by EPSRC grant EP/D073626/2 and NSF grant DMS-0602191. The second author was supported by an NSERC PDF, the ANR "Geometric, analytic and algorithmical aspects of groups", and a Fields Postdoctoral fellowship.

2 Dunwoody/Delzant-Potyagailo resolution

In this section we give a version of [DP01, §4]. Rather than work with free actions on simply connected spaces Delzant and Potyagailo use actions on simply connected triangular complexes where cell stabilizers are allowed to be members of an "elementary family".

Given a (reasonable) space X with a free G–action and a G–tree T there is always a G–equivariant map from X to T. If T is simplicial and the map is chosen reasonably well, preimages of midpoints of edges form a subset of X called a *pattern*, the connected components of which are two-sided *tracks*. Patterns were introduced by Dunwoody in [Dun85] to show that almost-finitely presented groups are accessible, and used in [DS99] to construct a JSJ decomposition for finitely presented groups over slender edge groups. If the action on X is not free there is in general no G–equivariant map $X \to T$. We coarsely follow [DP01] and construct a class of spaces such that if T is a (suitable) G–tree then there are G–equivariant maps $X \to T \cup \partial T$.

$\mathcal{H}$–complexes

All complexes in the sequel are at most two-dimensional. A CW–complex is *triangular* if all two-cells have at most three sides. Let H be a finitely generated group with a slender hierarchy $\mathcal{H}$. For $G < H$, an $\mathcal{H}$–complex for G is a connected two-dimensional triangular complex X with $H^1(X, \mathbb{Z}_2) = 0$, X/G compact, and slender or $\mathcal{H}$–elliptic cell stabilizers. An $\mathcal{H}$–complex is *nonelementary* if it contains a triangle, and is elementary if it doesn't.

We follow [DP01] in making a sequence of reductions to $\mathcal{H}$–complexes. If X contains a cell e which is attached to a single vertex of $X^{(0)}$ then collapse connected components $G \cdot e$ to points. Iterate until there are no loops or spheres. Suppose that X has a bigon (a 2–cell with two edges). Let $\mathcal{G}$ be the graph with a vertex for each bigon and an edge between two vertices if they're contained in the same bigon. Let Y be a subcomplex of X corresponding to a connected component of $\mathcal{G}$. The subcomplex Y has two vertices since X has no spheres. Collapse all bigons in Y to an edge h_Y, likewise for all translates of Y. Then $\mathrm{Fix}(h_Y)$ is contained in the stabilizer of the endpoints of h_Y, which are either slender or $\mathcal{H}$–elliptic. If $\mathrm{Stab}(h_Y)$ is not slender then collapse connected components of the union of translates of h_Y to points. Repeat.

Resolving actions on trees

This is essentially [DP01, Dessins 1 et 2], following [Dun85]. Denote the boundary of a tree T by ∂T. Suppose X is an $\mathcal{H}$–complex for $G < G' \in \mathcal{H}$ and let T be the tree associated to $\Delta_{G'}$. Cell stabilizers in X might not act elliptically in T, and there is therefore no G–equivariant map $X \to T$. By the definition of an $\mathcal{H}$–complex, a cell stabilizer in X fixes a point in $\hat{T} = T \cup \partial T$.

Let v be a representative of an orbit of vertices in X. Choose arbitrarily a point $\varphi(v) \in \mathrm{Fix}_T(\mathrm{Stab}(v)) \in \hat{T}$, provided that if $\mathrm{Stab}(v)$ is elliptic then $\varphi(v) \in T^0$. Note that since T is hyperbolic $\mathrm{Fix}_T(\mathrm{Stab}(c))$ is not empty for all cells c in X. For each $g \cdot v$ in the orbit of v, set $\varphi(g \cdot v) = g \cdot \varphi(v)$. Repeat over all orbits.

Let e be a representative of an orbit of edges of X, and let v and w be the endpoints of e. Suppose that $\mathrm{Stab}_X(e)$ inverts e and acts elliptically in T. Let m_e be the midpoint of e and let $a = \varphi(v)$, $b = \varphi(w)$, and choose $c \in \mathrm{Fix}_T(\mathrm{Stab}_X(e))$. Let r_a and r_b be the two rays in T connecting c to a and b, respectively. Let $\varphi(m_e)$ be the furthest point in $r_a \cap r_b$ from c if there is one. Otherwise $a = b \in \partial(T)$, and we set $\varphi(e) = a = b$. Then $\mathrm{Stab}(e)$ stabilizes the (possibly degenerate) arc in T connecting a and b. Map $[m_e, v]$ homeomorphically to the (possibly degenerate) interval connecting $\varphi(v)$ and $\varphi(m_e)$ in $\hat{T}$, extend equivariantly to all $[m_e, w] \subset e$, and then to all $g \cdot e$ in the orbit of e. Repeat over all edges.

The edges of a triangle t determine a (possibly degenerate) tripod in T. Map t to T equivariantly (t may have nontrivial stabilizer), and extend equivariantly to all translates of t. Repeat over all orbits of triangles. (As with edges which are inverted, some small amount of care must be taken since $\mathrm{Stab}(t)$ might not fix t.) See Figure 1.



Figure 1: Map on a typical triangle. The vertex in the lower left corner is sent to ∂T, which is indicated by the dotted lines. The center picture represents the triangle after crushing preimages of midpoints of edges, which introduces bigons and creates a new triangle.

Let G, X, and T be as above, and suppose $\varphi^{-1}(\partial T)$ doesn't contain any one-cells. Let $X^* = X \setminus \varphi^{-1}(\partial T)$, and let $\varphi\colon X^* \to T$ be the map obtained by restriction. For each edge e of T let m_e be the midpoint of e, and let Λ' be the one complex $\varphi^{-1}(\cup_e \{m_e\})$. Call a connected component λ of Λ' *essential* if both components of $X \setminus \lambda$ are unbounded and λ is not parallel to the link of a vertex, and let Λ be the union of all essential components of Λ'. Let X^*/Λ be the space obtained by collapsing each connected component of Λ to a point, and let X_T be the space obtained by reducing. The stabilizers of the resulting vertices are slender and there is a G–equivariant map $X^*/\Lambda \to X_T$.

Example 1 (Counterexample to [DP01, Lemma 4.10].). *It is erroneously claimed in [DP01, Lemma 4.10] that there is a map $X/G \to X_T/G$ with connected fibers, inducing an isomorphism on fundamental groups, hence that*

$$\mathrm{b}_1(X/G) \geq \sum_i \mathrm{b}_1(X_i/G_i) + \mathrm{b}_1(T/G),$$

where each X_i is a representatives of a cutpoint free component of X_T and G_i is its stabilizer. The following example is not of minimal complexity, but may perhaps be embedded in more interesting examples. See Figure 2. Consider a disk $X/\mathbb{Z}$ with one orbifold point, labeled $\mathbb{Z}$, such that the boundary of the disk defines a generator. The cyclic group $\mathbb{Z}$ acts on the line T, and applying the construction to the resolution $X \to \hat{T}$, where X is the (orbihedral) universal cover of the disk, recovers $\mathbb{Z}$ as the fundamental group of a space homotopy equivalent to a circle. Any continuous map from a disk to a circle, however, has nullhomotopic image, hence there is in general no natural map $X \to X_T$. There doesn't seem to be an obstruction to this phenomenon appearing in more complicated examples.

Lemma 2.1 (*c.f.* [DP01, Lemma 4.9]). $H^1(X^*/\Lambda, \mathbb{Z}_2) = 0$



Figure 2: [DP01, Lemma 4.10] violates the no retraction theorem.

Proof. Let Z be a connected component of X^*, let Y be its closure in X, and let W be the connected component of X^*/Λ corresponding to Z. Let c be a closed path in W. The quotient map $Z \to W$ is surjective on fundamental group, hence if c is a path in W then c is the image of a path d in Z. Since Y is 1-acyclic the path d is homologous to a sum of paths in $Z \subset Y$ which are individually homologous into links of vertices of Y, hence if each such path is homotopically trivial in W then $H^1(W, \mathbb{Z}_2) = 0$.

Let d be a path in $Z \subset Y$ which is homotopic into a vertex $v \in Y$. Then d is homotopic to a path d' in the link of v in Y. If $v \notin \varphi^{-1}(\partial T)$ then $v \in Z$ and d is clearly nullhomotopic in W. Let $e_1, \ldots, e_n$ be the edges in the link of v crossed by d', such that $d' = e_1 \cdots e_n$ as an edge path, and let $t_1, \ldots, t_n$ be the triangles in W with $e_i, \{v\} \subset t_i$. Since $\varphi(e_i) \neq \varphi(v)$ there is an edge e of T such that $\varphi^{-1}(m_e) \cap t_i$ is an arc a_i connecting e_i to e_{i+1} in t_i for all i. Then d' is homotopic to the path $a_1 \cdots a_n$, which is a path in Λ', thus d has nullhomotopic image in W, and W is 1-acyclic. □

Suppose now that $\varphi^{-1}(\partial T)$ contains one-cell. We are concerned with the possibility that $\mathrm{Stab}(\varphi^{-1}(p))$, for $p \in \partial T$, is small, not slender, and corresponds to a cutpoint in X', i.e., corresponds to a splitting over a small but not slender edge group. Let X' be the complex obtained by collapsing each connected component of $\varphi^{-1}(\partial T)$ to a point and reducing. The following is immediate.

Lemma 2.2. *Vertex stabilizers in X' are either vertex stabilizers from X, or are either slender or small, f.g., and act parabolically on T. Edge stabilizers in X' are slender.*

Since X' potentially has small but not slender vertex stabilizers it is *not*, in general, an $\mathcal{H}$–complex.

Proof. Let v be a vertex in X'. Denote by $\mathrm{Stab}(v)$ by G_v. If v is a vertex from X then clearly G_v is a vertex stabilizer from X. Suppose v corresponds to a connected component V of $\varphi^{-1}(p)$, with $p \in \partial T$. Clearly $\mathrm{Stab}(V) = G_v$. If $gV \cap V \neq \emptyset$ then $g \cdot p = p$ and therefore $gV = V$, hence $V/G_v \to X/G$ is an embedding and G_v fixes p. Since X/G is compact G_v acts cocompactly on V, and since V has slender cell stabilizers G_v is finitely generated. Let $T' \subset T$ be the minimal subtree stabilized by G_v, and consider the quotient T'/G_v. Since G_v fixes an end in T, it fixes an end in T', and T'/G_v is therefore an ascending HNN extension with slender edge groups, hence is either small or slender, and if small, it acts parabolically on T.

Let e be an edge in X' and let G_e be its stabilizer. Then G_e fixes the endpoints v and w of e. Let V and W be the preimages of v and w in X. Then G_e stabilizes both V and W. If V and W are connected components of preimages of points in ∂T then G_e fixes a pair of distinct points in ∂T, hence G_e is slender. If V is not a connected component of a preimage of any point in ∂T then V is a vertex, and G_e stabilizes a half-line in T, hence is slender in this case as well. □

The complexity of an $\mathcal{H}$–complex X for G is the number of triangles in X/G, and is denoted unambiguously by $\operatorname{covol}(X)$.

Small edge stabilizers

It is tempting to try to extend the technique to groups acting on trees with small edge stabilizers. There is no trouble with this when edge stabilizers in X act elliptically in T, but this isn't always the case.

Example 2. *Consider the* HNN *extension G, with associated graphs of groups decomposition Δ, of $\mathbb{Z}^3$ with edge group $\mathbb{Z}^3$, determined by the following two matrices:*

$$\begin{pmatrix} n & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & m \end{pmatrix}$$

Set $A = \mathrm{BS}(1,n)$, $B = \mathrm{BS}(1,m)$, and let t and t' be the stable letters associated to A and B. Then G is a quotient of

$$(A \oplus \langle c\rangle) *_{\langle c'', t''\rangle^{ab}} (B \oplus \langle c'\rangle)$$

with edge maps $c'' \mapsto c, t'' \mapsto t^{-1}$ and $c'' \mapsto c', t'' \mapsto t'$. Note that $t'' \mapsto t^{-1}$. Let $C = \langle c'', t''\rangle^{ab}$. Furthermore, G acts on a simply connected space with vertex and edge stabilizers conjugate to $A \oplus \langle c\rangle$, $B \oplus \langle c'\rangle$, and C. Then A and B fix distinct points in the boundary of the tree T associated to Δ and C acts hyperbolically in T, hence there is no place to map an edge stabilized by C.

Examples like this might occur in nature.

3 Products of trees

Let G be a group and let T and T' be a pair of G–trees with $T/G = \Delta$ and $T'/G = \Omega$. Then G acts diagonally on the product $T \times T'$ with quotient a square complex. Following [FP06] we think of S as a complex of groups. Let

$S \subset T \times T'$ be a simply connected G–invariant subset, and suppose that point preimages under π_T are connected. For v a vertex of Δ let $\widetilde{v}$ be a lift of v to T. Then $G_v \cong \mathrm{Stab}(\widetilde{v})$ acts on the tree $\pi_T^{-1}(\widetilde{v})$ and $\pi_T^{-1}(\widetilde{v})/G_v$ is a graph of groups decomposition of G_v. Similarly, if m_e is a midpoint of an edge of Δ then $\pi_T^{-1}(\widetilde{m}_e)/G_e$ is a graph of groups decomposition of G_e.

Theorem 3.1 (*c.f.* [Gui05, Théorème principal, Corollaire 8.2])**.** *There is a connected simply connected G–equivariant square complex $S \subset T \times T'$ of minimal covolume such that the projections $S \to T'$ and $S \to T$ have connected point preimages.*

Moreover, if G and all vertex and edge stabilizers are finitely generated and T' is a minimal invariant subtree then S/G is compact.

It is necessary in what follows that we allow T to not be a minimal invariant tree.

Let $S_{\Delta,\Omega} = S/G$, and denote the projections $S_{\Delta,\Omega} \to \Delta, \Omega$ by π_Δ and π_Ω. We have that $S_{\Delta,\Omega}$ is finite, and if v is a vertex in Δ, then $\pi_\Delta^{-1}(v)$ is a graph of groups decomposition of G_v, and if m is a midpoint of an edge of Δ then $\pi_\Delta^{-1}(m)$ is a graph of groups decomposition of G_e, likewise for Ω.

4 $\mathcal{H}$–structures

There is no natural way to construct an $\mathcal{H}$–complex for each group L in a hierarchy $\mathcal{H}$. To get around this difficulty we define an $\mathcal{H}$–structure, which is a combination of a hierarchy (distinct from $\mathcal{H}$!) and a collection $\mathcal{H}$–complexes for terminal groups in the hierarchy. We associate to each group L in $\mathcal{H}$ an $\mathcal{H}$–structure $\mathcal{X}_L$ and show in §5 that for groups sufficiently far down the hierarchy the $\mathcal{H}$–structures are essentially hierarchies whose terminal vertex groups are $\mathcal{H}$–elliptic or slender.

Definition 4.1. Let $\mathcal{H}$ be a hierarchy for a group H. An *$\mathcal{H}$–structure* of a group $L < H$ is a finite hierarchy over slender or small edge groups equipped with an action, for each terminal vertex group V of $\mathcal{X}_L$, of V on an $\mathcal{H}$–complex X_V. The complexity $\mathrm{covol}(\mathcal{X}_L)$ is the total number of orbits of triangles over all X_V.

We indicate that an $\mathcal{H}$–structure for L is over slender edge groups by denoting it by $\mathcal{X}_L$, and if an $\mathcal{H}$–structure for L possibly has small edge groups then we denote it by $\mathcal{Y}_L$. If $L \in \mathcal{H}$ has has an $\mathcal{H}$–structure $\mathcal{Y}_L$ we require that all non-slender small edge groups in $\mathcal{Y}_L$ act parabolically in T_{Δ_L}.

The *height* of an $\mathcal{H}$–structure on L is the number of levels in $\mathcal{X}_L$, and is denoted by $\mathrm{height}(\mathcal{X}_L)$. We denote graphs of groups decompositions in $\mathcal{H}$–structures by Ω, i.e., if $L' \in \mathcal{X}_L$ then the graph of groups decomposition of L' will be denoted $\Omega_{L'}$.

Given $G \in \mathcal{H}$ and an $\mathcal{H}$–structure on G we would like to find, for each vertex group Z of Δ_G, an $\mathcal{H}$–structure $\mathcal{X}_Z$. For each terminal group B in $\mathcal{X}_G$ acting on a nonelementary $\mathcal{H}$–complex X_B we first use §2 to replace B by an $\mathcal{H}$–structure $\mathcal{X}_B$ so the terminal groups of $\mathcal{X}_B$ are elliptic in T.

Resolving the action of G on T

Let X be a triangular complex with a G action. Define a tree T_X as follows. First, let T be the tree whose vertices are the maximal cutpoint free components of X (maximal connected subcomplexes which aren't separated by a vertex) and the cutpoints of X. If v is a cutpoint and Y is a cutpoint free component such that $v \in Y$ then join $[v]$ and $[Y]$ by an edge $[l]$, where l is the connected component of the link of v contained in Y. Collapse edges with non-small stabilizers to obtain T_X.

Lemma 4.2. *Let $\mathcal{X}_G$ be an $\mathcal{H}$–structure for $G \in \mathcal{H}$. There are $\mathcal{H}$–structures $\mathcal{X}_Z$, $Z \in \Delta_G$ such that*

$$\sum_{Z \in \Delta_G} \operatorname{covol}(\mathcal{X}_Z) \leq \operatorname{covol}(\mathcal{X}_G)$$

and if, for each terminal vertex group B of $\mathcal{X}_G$, X_B is a point, then

$$\operatorname{height}(\mathcal{X}_Z) \leq \operatorname{height}(\mathcal{X}_G)$$

Proof. Let $\mathcal{X}_G$ be an $\mathcal{H}$–structure on G, and suppose G acts on a slender G–tree T with quotient Δ_G. Suppose $\mathcal{X}_G$ has a nonelementary terminal vertex group B acting on an $\mathcal{H}$–complex X_B. Let X' be the complex associated to B and T provided by Lemma 2.2. Let Ω_B be the graphs of groups decomposition $T_{X'}/B$, and for each vertex v of Ω let X_v be the subcomplex of X' stabilized by B_v. There is a natural B_v–map $X_v \to T$ obtained by restriction, with X_v^* connected. Let Y_v be the $\mathcal{H}$–complex $(X_v)_T$. Now let $\Omega_{B_v} = T_{Y_v}/B_v$, and for each vertex w of T_{Y_v}/B_v let $B_{v,w}$ act on the subcomplex of Y_v corresponding to a lift of w. Repeat over all nonelementary terminal vertex groups B of G to obtain an $\mathcal{H}$–structure $\mathcal{Y}_G$.

If there are no nonelementary terminal vertex groups set $\mathcal{Y}_G = \mathcal{X}_G$. Each terminal vertex group B in the resulting decomposition is either $\mathcal{H}$–elliptic or slender.

Let Ω_L be the decomposition of L for $L \in \mathcal{Y}_G$. For each vertex group L' of Ω_L define inductively $\Delta_{L'} = \pi_\Omega^{-1}(L') \subset \mathcal{S}_{\Delta_L,\Omega_L}$. See Figure 3 We have constructed a hierarchy of square complexes modeled on $\mathcal{Y}_G$: If L' is a (nonterminal) vertex group of Ω_L there is a natural projection

$$S_{\Delta_{L'},\Omega_{L'}} \to \Delta_{L'} = \pi_\Omega^{-1}(L') \subset S_{\Delta_L,\Omega_L}$$

There is a natural map Π from this hierarchy of square complexes to Δ_L, and if Z is a vertex group of Δ_L then $\mathcal{X}_Z = \Pi^{-1}(Z)$ is almost an $\mathcal{H}$–structure on Z. Let W be a terminal vertex group of $\mathcal{Y}_G$.

Suppose W is a nonelementary terminal vertex group of $\mathcal{Y}_G$. Since X_W is nonelementary then Δ_W is a finite tree representing the trivial graph of groups decomposition of W, and for each vertex group V of Δ_W in $\mathcal{X}_Z$, if V is slender let X_V be a point, and if $V = W < W$ let $X_V = X_W$. □



Figure 3: A piece of the hierarchy of square complexes associated to $\mathcal{Y}_G$.

Hierarchy of $\mathcal{H}$–structures

Let H be a finitely generated group, $\mathcal{H}$ a slender hyperbolic hierarchy for H, and suppose H is $\mathcal{H}$–almost finitely presented. Let $\mathcal{X}_H$ be the trivial $\mathcal{H}$–structure with trivial graph of groups decomposition, and let X_H be any $\mathcal{H}$–complex for H.

Suppose for $L \in \mathcal{H}$, that $\mathcal{X}_L$ has been defined. For Δ_L and Z a vertex group of Δ_L, let $\mathcal{X}_Z$ be the $\mathcal{H}$–structure on Z constructed in the previous subsection. Let $B_{L,1}, \dots, B_{L,n_L}$ be the terminal vertex groups acting on nonelementary $\mathcal{H}$–complexes $X_{B_{L,i}}$. Let $L_1, \dots, L_k$ be the descendants of L. Then each $X_{B_{L_j,k}}$ is obtained from some $X_{B_{L,i(k)}}$ by resolving the action of $B_{L,i(k)}$ on T_{Δ_L}. We call the collection of $B_{L_j,i(k)}$ such that $i(k) = i$ the *descendants* of B_i^L. Since X_H has finitely many triangles, for all but finitely many L, each $B_{L,i}$ has exactly one descendant $B_{L_j,i}$ and $\operatorname{covol}(X_{B_{L,i}}) = \operatorname{covol}(X_{B_{L_j,i}})$. We have:

Lemma 4.3 (*c.f.* [DP01, p. 627])**.**

$$\operatorname{covol}(X_{B_{L,i}}) \geq \sum_{k|i(k)=i} \operatorname{covol}(X_{B_{L_j,k}})$$

and for all but finitely many $L \in \mathcal{H}$, the sum on the right is over one element and the inequality is an equality. There is some N_0 so that for $i \geq N_0$ this is the case.

In what follows we assume $i \geq N_0$.

5 Nonelementary complexes converge to trees

The aim of this section is to replace, for groups sufficiently far down the hierarchy $\mathcal{H}$, each $\mathcal{H}$–structure $\mathcal{X}_L$ with an $\mathcal{H}$–structure with no triangles.

Consider the finite collection of infinite sequences of terminal vertex groups $\{G^p_{N_0} > G^p_{N_0+1} > G^p_{N_0+2} > \dots\}$ such that $G^p_i \in \mathcal{X}_{L(p,i)}$, $L(p,i) \in \mathcal{H}^i$, is terminal, acts on a nonelementary $\mathcal{H}$–complex $X^p_{G^p_i}$, is the only descendant of G^p_{i-1}, such that $\operatorname{covol}(X^p_{G^p_i}) = \operatorname{covol}(X^p_{G^p_{i+1}})$. To simplify notation we drop the

'p's and denote G_i^p by G_i and $X^p_{G_i^p}$ by X_i. Note that if v is a vertex in X_i with slender stabilizer then the link l of v has exactly one connected component and $\mathrm{Stab}(l) = \mathrm{Stab}(v)$.

Lemma 5.1. *Edge and face (but not necessarily vertex!) stabilizers in X_i are $\mathcal{H}$–elliptic.*

Proof. Otherwise for some i, $\varphi_i^{-1}(\partial T_i)$ contains a 1–cell which, when collapsed, decreases the number of orbits of triangles, hence $\mathrm{covol}(X_i) > \mathrm{covol}(X_{i+1})$. $\square$

Let Δ_i be the decomposition G_i inherits from $\Delta_{L(p,i)}$, and let T_i be the associated tree. We argue that for sufficiently large i, X_i can be replaced by a graph of groups such that each vertex group acts on a tree with $\mathcal{H}$–elliptic or slender vertex stabilizers.

Let $\mathcal{L}_i$ be the collection of orbits of connected components of links of vertices of X_i, and denote the orbit of a link l by $[l]$. Say that $[l] \in \mathcal{L}_i$ *dies* if $\mathrm{Stab}(l)$ acts hyperbolically in T_i, otherwise $[l]$ *survives.* Let $\mathcal{L}_i^s$ be the collection of orbits of connected components of links of vertices which survive, and let $\mathcal{I}_i$ be the collection of orbits of links of vertices which survive forever. There is a natural map $\iota_i \colon \mathcal{I}_i \to \mathcal{I}_{i+1}$, and since $\mathrm{covol}(X_{i+1}) \le \mathrm{covol}(X_i) \le \mathrm{covol}(X_H)$, eventually ι_i is bijective. Clearly the links which survive forever have $\mathcal{H}$–elliptic stabilizers, and we call a link l with $\mathcal{H}$–elliptic stabilizer $\mathcal{H}$–elliptic.

Let $\mathcal{V}_i$ be the collection of orbits of vertices v such that v has an $\mathcal{H}$–elliptic components in its link. Since the number of orbits of connected components of links which survive forever is constant, $|\mathcal{V}_i|$ is non-decreasing in i, and is eventually constant. Furthermore, if a component of the link of v is not slender then *all* components of the link of v have non-slender stabilizer, and if the link of v has an $\mathcal{H}$–elliptic component l and $\mathrm{Stab}(l)$ is slender then l is the only component of $\mathrm{lk}(v)$.

Let l be an $\mathcal{H}$–elliptic component of a link. Then

$$|l/\operatorname{Stab}(l)| \le |\iota_i(l)/\operatorname{Stab}(\iota_i(l))|$$

For sufficiently large i this number stabilizes as well, giving a bijection $\mathcal{V}_i \to \mathcal{V}_{i+1}$ and, for each vertex v with an $\mathcal{H}$–elliptic link component, a $\mathrm{Stab}(v)$–equivariant isomorphism of links $\mathrm{lk}_{X_i}(v) \to \mathrm{lk}_{X_{i+1}}(\varphi_i(v))$. We assume below that $N_1 \ge N_0$ is chosen large enough to arrange all of the above, over all sequences $\{G_i^p\}$, for $i \ge N_1$.

Pairs of triangles

For $i \ge N_1$, X_i^* is connected and $\varphi_i \colon X_i^* \to (X_i)_{T_i}$ induces bijections on orbits of triangles and stars of vertices with $\mathcal{H}$–elliptic components in their links.

Let triangles(X_i) be the set of triangles in X_i, and let $\tau_i\colon$ triangles$(X_i) \to$ triangles$((X_i)_{T_i})$ be the induced map. A pair of triangles is a triple $P = (t, t', e)$ where $t, t' \in$ triangles(X_i) are triangles meeting in the edge e. Denote the collection of pairs of triangles in X_i by triangles$^2(X_i)$. The map φ_i *separates* two triangles t and t' if $\tau_i(t)$ and $\tau_i(t')$ lie in different cutpoint free components of $(X_i)_{T_i}$ (Each track λ separates X_i into two components.). See Figure 4. If φ_i doesn't separate P then $[P] \in \text{triangles}^2(X_i)/G_i$ descends to an equivalence class of pairs $\tau_i([P])$ of triangles in $\text{triangles}^2(X_{i+1})/G_{i+1}$. Similarly, say that φ_{i+1} doesn't separate P if φ_i doesn't separate P and if $P' \in \tau_i([P])$ then φ_{i+1} doesn't separate P_i. Likewise for φ_k for $k = i+2, \ldots$. Let $\text{spairs}(X_i)$ be the collection of (stable) pairs of triangles which are not eventually separated by any φ_j.



Figure 4: An adjacent pair of triangles separated by T.

The following lemma is obvious from the definitions.

Lemma 5.2. *If* $[(s,t,e)], [(s,t',e)] \in \text{spairs}(X_i)$ *then* $[(t,t',e)] \in \text{spairs}(X_i)$. *If* $[(t_0,t_1,e_0)], [(t_1,t_2,e_1)], \ldots, [(t_{n-1},t_n,e_{n-1})] \in \text{spairs}(X_i)$ *and* t_0 *and* t_n *share an edge* e *then* $[(t_0,t_n,e)] \in \text{spairs}(X_i)$.

Let $\sim_i$ be the equivalence relation on $\text{triangles}(X_i)$ generated by $s \sim_i t$ if $[(s,t,e)] \in \text{spairs}(X_i)/G_i$. There are induced (injective) maps

$$\sigma_{i,j} \colon \text{spairs}(X_i)/G_i \to \text{spairs}(X_j)/G_j$$

which push forward the equivalence relations $\sim_i$ to equivalence relations $\sigma_{i,j}(\sim_i)$ on $\text{triangles}(X_j)$ which are compatible with, and perhaps finer than, $\sim_j$. Let $P_i^1, \ldots, P_i^{n_i}$ be subcomplexes of X_i which represent orbits of unions of $\sim_i$ equivalence classes. Then $\cup_j (G_i/\operatorname{Stab}(P_i^j))P_i^j$ contains all triangles in X_i, and by Lemma 5.2, if gP_i^j and $hP_i^{j'}$ overlap in a triangle then $j = j'$ and $h^{-1}g \in \operatorname{Stab}(P_i^j)$. Let

$$D_i := \left(n_j, \sum_j \chi((P_i^j \setminus (P_i^j)^0)/\operatorname{Stab}(P_i^j)) \right)$$

Let P_j^i be the union of triangles in a $\sim_i$ equivalence class, and let γ_j^i be the graph obtained by restricting to edges of the barycentric subdivision of P_j^i which don't meet vertices of P_j^i. Each term in the sum in the second term of D_i is simply the Euler characteristic of $\gamma_j^i/\operatorname{Stab}(P_j^i)$. It is possible to realize the γ_j^i disjointly and equivariantly in X_i as a graph which "guides" the pattern Λ_i around X_i.

Lemma 5.3. *If $\sigma_{i,j}(\sim_i)$ is strictly finer than $\sim_j$ then $D_i > D_j$.*

Proof. If $\sigma_{i,j}(\sim_i)$ is strictly finer than $\sim_j$ then there is an additional stable pair of triangles which either joins two components, reducing the first coordinate, or joins a component and a translate of itself, which reduces the second. $\square$

The complexity is controlled from below by $\operatorname{covol}(X_i)$:

$$D_i \geq (1, 1 - 2\operatorname{covol}(X_H))$$

and there is therefore some first index $M_p \geq N_1$ (recall we are working in the branch $G_i^p = G_i$) such that $D_{M_p} = D_j$ for $j \geq M_p$.

An *unstable* edge is an edge e such that there are triangles t, t' so that $[(t, t', e)] \notin \operatorname{spairs}(X_i)$. Let $W \subset X_i$ be the collection of unstable edges in X_i.

A *cone* C is a triangulated disk with exactly one interior vertex. A cone in a triangular complex X is a combinatorial map $\gamma \colon C \to X$ which maps triangles to triangles. A cone $\gamma \colon C \to X$ is *simple* if the associated path in the link of the image of the cone point is simple.

Let $C \to X$ be a cone in X and let C^* be the space obtained by removing vertices of C which are mapped to $\varphi^{-1}(\partial T)$. Let λ be the preimage of Λ in C^*. The map $C^* \to X^*/\Lambda$, induces a natural map $C' \to X_T$, where C' is the space obtained by collapsing each connected component of λ in C to a point, followed by collapsing bigons to edges.

Let $C \to X$ be a cone and suppose $X \to T$ is as above. Let c be the cone point in C, and let s be the outermost component of λ encircling the cone point, if there is one, otherwise let $s = c$. The *push-forward* C'' of C to X_T is the cone obtained from C' by taking all triangles in C' containing the image of s. See Figure 5.



Figure 5: Constructing the push-forward C'' of C.

Lemma 5.4. *Suppose $C_i \to X_i$ is a simple cone, and that there are two triangles t and t' in the image of C_i such that $t \not\sim_i t'$. Then $i < M_p$.*

Proof. Suppose $i \geq M_p$. Let $C_j \to X_j$ be the push-forward of C_i to X_j. Since there are triangles t and t' such that $t \not\sim_i t'$ then for some $j > i$, $|C_j| < |C_i|$. Let j be the first index such that $|C_j| = |C_{j'}|$ for $j' \geq j$. Then all triangles in the image of C_j are $\sim_j$ equivalent. Let $t_1, \ldots, t_n$ be the triangles in the image of C_j, indexed so that $[(t_k, t_{k+1}, e_k)] \in \operatorname{spairs}(X_j)/G_j$. Let $\tilde{t}_k$ be the triangle in the image of C_i in X_i corresponding to t_k. Then since $i \geq M_p$ there are edges $\tilde{e}_k$ in X_i such that $\big[(\tilde{t}_k, \tilde{t}_{k+1}, \tilde{e}_k)\big] \in \operatorname{spairs}(X_i)/G_i$, but this implies that the cone $C_i \to X_i$ was not simple. $\square$

Lemma 5.5. *Suppose $i > M_p$. If e is unstable, i.e., there are triangles t and t' such that $[(t, t', e)] \notin \operatorname{spairs}(X_i)$, then e separates X_i, with $t \setminus e$ and $t' \setminus e$ lying in different components of $X_i \setminus e$.*

Note that there may be edges which separate X_i, but which don't count as unstable edges.



Figure 6: The homology h from Lemma 5.5, which we portray as a disk, even though it need not be planar. The edges with arrows are mapped to e.

Proof of Lemma 5.5. Let a and b be the vertices of t and t', respectively, distinct from the endpoints v and w of e. Suppose that e doesn't separate X_i into at least two components, with $t \setminus e$ lying in one and $t' \setminus e$ lying in another. Then there is an edge path $p\colon I \to X_i$ of a subdivided interval such that $p(0) = a$ and $p(1) = b$ such that $p^{-1}(e) = \emptyset$. Let f and g be the oriented edges of t and t' connecting a to v and v to b, respectively. Let $h'\colon D' \to X_i$ be a combinatorial map of a triangulated surface D' representing a homology between the edge paths gf and p, and let $h\colon D \to X_i$ be the combinatorial map of a surface obtained by attaching two triangles representing $t \cup_e t'$ to D'. See Figure 6.

Without loss, by perhaps changing p and h, we may assume that the union of edges of D which are mapped to e does not separate D, and we may assume that D has the form illustrated in Figure 6. The path p may be divided into subpaths $p_0, \ldots, p_{n-1}$ such that p_j connects the apex of a triangle t_j to the apex of a triangle t_{j+1}, and such that the side of t_j is mapped to e by h. Furthermore, by identifying the edges labeled e in the sequence of triangles determined by t_j, p_j, and t_{j+1}, we obtain a cone in the link of one of v or w. Then either $t_j = t_{j+1}$ or there is a simple cone $C \to X_i$ containing t_j and t_{j+1}, hence $t_j \sim_i t_{j+1}$ for all j, hence $t \sim_i t'$ contrary to hypothesis. $\square$

Lemma 5.6. *Suppose* $[(t,t',e)] \notin \mathrm{spairs}(X_i)$. *Then neither stabilizer of a link of a vertex corresponding to* e *is* $\mathcal{H}$*–elliptic.*

Proof. Suppose not. Let v and w be the vertices joined by e, such that the component l of the link of v corresponding to e is $\mathcal{H}$–elliptic. Then the link of v in X_j, for some $j > i$, has strictly fewer orbits of edges than l, contradicting our choice of N_1. $\square$

Replacing X_i by a tree of planes

Fix some $i \geq N_2 > \max_p\{M_p\}$, let W be the union of unstable edges in X_i, and let $\{P_\alpha\}$ be the collection of maximal connected subcomplexes which aren't eventually separated by any φ_j.

Lemma 5.7. $\mathrm{Stab}(P_\alpha)$ *is* $\mathcal{H}$*–elliptic.*

Proof. Clearly P_α is a union of $\sim_i$ equivalence classes. Let $P \neq P' \subset P_\alpha$ be unions of $\sim_i$ equivalence classes with nonempty intersection. If P and P' share an edge then there are triangles $t \subset P$ and $t' \subset P'$ which share an edge e, but then $[(t,t',e)] \in \mathrm{spairs}(X_i)$, hence $P = P'$. Suppose P and P' meet in a vertex v. If λ is a connected component of Λ_i and separates a pair of triangles $t \subset P$ and $t' \subset P'$ then λ is parallel to the link of v, but then a component of $\mathrm{lk}(v)$ has stabilizer which doesn't survive forever, contradicting our choice of N_1. $\square$

For each P_α introduce a vertex v_α and let P'_α be the cone on $W \cap P_\alpha$ with cone point v_α, and define a new $\mathcal{H}$–complex X'_i for G_i by replacing each P_α by P'_α.



Figure 7: The link of a vertex in $W \subset X'_i$, doesn't contain a loop, otherwise complexity decreases further down the hierarchy.

Lemma 5.8. *Links of vertices in* $W \subset X'_i$ *are not* $\mathcal{H}$*–elliptic, and are lines with finite trees attached.*

See Figure 7.

Proof. The first part of the lemma is obvious by our choice of N_1. By construction, if $v \in W$ then $\mathrm{lk}(v)$ is connected. We first show that the link of a vertex

$v \in W$ is a tree. Suppose it isn't. Then there is an simple loop c in $\mathrm{lk}(v)$. Let $e_0, \ldots, e_n$ be the cyclically ordered unstable edges in $W \cap \mathrm{lk}(v)$ corresponding to c. For each pair e_j, e_{j+1} let β_j be the index such that there is a pair of triangles $t_j, t_j' \subset P'_{\beta_j}$ such that $e_j \subset t_j$, $e_{j+1} \subset t_j'$, and t_j and t_j' share the edge connecting v_{β_j} to v. Since c is simple, $\beta_j \neq \beta_{j'}$ if $j \neq j'$. There is a simple chain of triangles $t^j_1, \ldots, t^j_{m_j}$ connecting e_j and e_{j+1} in $\mathrm{lk}_{P_{\beta_j}}(v)$ and

$$t^0_1, \ldots, t^0_{m_0}, \ldots, t^n_1, \ldots, t^n_{m_n}$$

gives a simple cone in $\mathrm{lk}_{X_i}(v)$, but $t^j_k \not\sim_i t^{j'}_{k'}$ for some j, j', k, k', $j \neq j'$ or $j =' $ and $k \neq k'$, contradicting Lemma 5.4.

Let l be a link of a vertex v in X_i' corresponding to a non-$\mathcal{H}$–elliptic link in X_i. Then l is a tree with slender stabilizer and the vertices of l correspond either to unstable edges adjacent to v or subgraphs of $\mathrm{lk}_{X_i}(v)$ which *aren't* separated by unstable edges.

Suppose $\mathrm{Stab}(l)$ acts on l with a global fixed point. Then either $\mathrm{Stab}(l)$ stabilizes an unstable edge in X_i, contradicting Lemma 5.1, or, for some α, stabilizes $\mathrm{lk}_{P_\alpha}(v) \subset \mathrm{lk}_{X_i}(v)$, which isn't separated by an unstable edge. But then $\mathrm{Stab}(l) < \mathrm{Stab}(P_\alpha)$ is $\mathcal{H}$–elliptic, contradicting 5.6.

Since $\mathrm{Stab}(l)$ is slender and doesn't fix a point in l it has an axis $A \subset l$, however $\mathrm{Stab}(l)$ acts cocompactly on l hence is obtained from A by attaching finite trees. $\square$

Thus every non-$\mathcal{H}$–elliptic link l in X_i' is an infinite line with finite trees attached. Each valence one vertex of l corresponds to a free face in X_i'. If f is a free face of a triangle t, equivariantly remove the interiors of f and t, and if there are any free faces repeat, decreasing the covolume until there are no free faces, to obtain X_i''. The link of each vertex in $W^{(0)} \subset X_i''$ is either a line or a collection of points (Faces opposite a vertex in W corresponding to an edge in $A \subset l$ may be free, and collapsing one of these turns the axis $A \subset l$ into a collection of finite trees. Further collapsing turns l into a collection of points.), and every edge adjacent to one triangle is adjacent to exactly two triangles, hence each component of the link of v_α in X_i'' is either a point, a line, or a circle. Edge stabilizers in X_i'' are contained in stabilizers of non-$\mathcal{H}$–elliptic links, hence are slender, thus each component of a link of a vertex in X_i'' has slender stabilizer. Refine the $\mathcal{H}$–structure $\mathcal{X}_{L(p,i)}$, by giving $G_i = G_i^p \in \mathcal{X}_{L(p,i)}$ the graph of groups decomposition $T_{X_i''}/G_i$, and for each vertex w of $T_{X_i''}/G_i$, assign the group $G_{i,w} = \mathrm{Stab}(\widetilde{w})$, which is either $\mathcal{H}$–elliptic or slender, in which case we set $X_{G_{i,w}}$ to be a point, or the stabilizer of a cutpoint free component Y of X_i'', in which case we set $X_{G_{i,w}} = Y$.

Lemma 5.9. *If $X_{G_{i,w}}$ is a cutpoint free component of X_i'' then $G_{i,w}$ acts on a tree with slender vertex stabilizers.*

Proof. Let Y be the nontrivial cutpoint free component of X_i'' stabilized by $G_{i,w}$, and let $W' = W \cap Y$ be the collection of unstable edges contained in Y, each of which bounds two triangles. Let v be a vertex in Y corresponding to some v_α. If v is connected to finitely many unstable edges in Y (say n), replace the star of v by an n-gon to obtain a polygonal complex Y'. Then every edge of Y' separates Y'. (The one-skeleton of Y' is in fact a subgraph of the Farey graph.) Let T be the dual tree, i.e., the tree with a vertex for each edge or polygon and an edge between an edge and a polygon containing it. Then $G_{i,w}$ acts on T with slender vertex stabilizers. $\square$

Proof of Theorem 1.3. Replace the $G_{i,w} = G^{p}_{i,w}$–complex Y by the tree T given by Lemma 5.9, then give $G_{i,w}$ the graph of groups decomposition $T/G_{i,w}$. $\square$

Remark 5.10. *Vertex groups in the* JSJ *decomposition of* G_i *are either slender, slender-by-orbifold, or* $\mathcal{H}$*–elliptic.*

6 JSJ hierarchy

As it is rather long and technical, we will not restate the definition of the JSJ decomposition of a finitely presented group over slender edge groups here, and instead refer the reader to [FP06, Theorem 5.13] and [DS99]. We need the following from [FP06].

Theorem ([FP06, Theorem 5.15])**.** *Let* G *be a finitely presented group, and* Γ *a graph decomposition we obtain in [FP06, Theorem 5.13]* [Note: Γ is the slender JSJ.]. *Let* $G = A *_C B$, $A*_C$ *be a splitting along a slender group* C*, and* T_C *its BassSerre tree.*

1. *If the group* C *is elliptic with respect to any minimal splitting of* G *along a slender group, then all vertex groups of are elliptic on TC .*
2. *Suppose the group* C *is hyperbolic with respect to some minimal splitting of* G *along a slender group. Then*
   - *(a) All non-enclosing vertex groups of* Γ *are elliptic on* T_C*.*
   - *(b) For each enclosing vertex group,* V*, of* Γ*, there is a graph de- composition of* V*,* $\mathcal{V}$*, whose edge groups are in conjugates of* C*, which we can substitute for* V *in* Γ *such that if we substitute for all enclosing vertex groups of* Γ *then all vertex groups of the resulting refinement of* Γ *are elliptic on* T_C*.*

Proof of Corollary 1.4. Let $L \in \mathcal{H}^n$ for some $n > N$, where N is as in Theorem 1.3. Groups in $\mathcal{H}$ are either slender-by-orbifold, hence have JSJ decompositions which are graphs of slender groups over slender edge groups, or are elliptic in the top level of the $\mathcal{H}$–structure of their parents, hence if $K < K'$ is a vertex group of the JSJ of K' and is not a graph of slender groups then we may assume height$(\mathcal{X}_K) <$ height$(\mathcal{X}'_K)$, hence $\mathcal{H}_L$ is finite and has terminal leaves which are either slender or are the non-slender terminal leaves of $\mathcal{X}_L$. $\square$

References

[BF91a] Mladen Bestvina and Mark Feighn, *Bounding the complexity of simplicial group actions on trees*, Invent. Math. **103** (1991), no. 3, 449–469. MR MR1091614 (92c:20044)

[BF91b] ______, *A counterexample to generalized accessibility*, Arboreal group theory (Berkeley, CA, 1988), Math. Sci. Res. Inst. Publ., vol. 19, Springer, New York, 1991, pp. 133–141. MR 1105331 (92g:20038)

[DJ98] M. J. Dunwoody and J. M. Jones, *A group with strange decomposition properties*, J. Group Theory **1** (1998), no. 3, 301–305. MR 1633200 (99d:20036)

[DP01] Thomas Delzant and Leonid Potyagailo, *Accessibilité hiérarchique des groupes de présentation finie*, Topology **40** (2001), no. 3, 617–629. MR MR1838998 (2002d:20063)

[DS99] M. J. Dunwoody and M. E. Sageev, *JSJ-splittings for finitely presented groups over slender groups*, Invent. Math. **135** (1999), no. 1, 25–44. MR MR1664694 (2000b:20050)

[Dun85] M. J. Dunwoody, *The accessibility of finitely presented groups*, Invent. Math. **81** (1985), no. 3, 449–457. MR 807066 (87d:20037)

[Dun93] ______, *An inaccessible group*, Geometric group theory, Vol. 1 (Sussex, 1991), London Math. Soc. Lecture Note Ser., vol. 181, Cambridge Univ. Press, Cambridge, 1993, pp. 75–78. MR 1238516 (94i:20067)

[FP06] K. Fujiwara and P. Papasoglu, *JSJ-decompositions of finitely presented groups and complexes of groups*, Geom. Funct. Anal. **16** (2006), no. 1, 70–125. MR MR2221253 (2007c:20100)

[Gui05] Vincent Guirardel, *Cœur et nombre d'intersection pour les actions de groupes sur les arbres*, Ann. Sci. École Norm. Sup. (4) **38** (2005), no. 6, 847–888. MR 2216833 (2007e:20055)

email: `lars@d503.net`

email: `nicholas.touikan@gmail.com`